

KISELEV, P.N.; OKULOV, N.M. [deceased]

Disorders in the absorption of various substances from the  
gastrointestinal tract in radiation sickness. Vop.radiobiol.  
2:199-212 '57. (MIRA 12:6)

1. Sotrudniki Tsentral'nogo nauchno-issledovatel'skogo rentgeno-  
radiologicheskogo instituta Ministerstva zdravookhraneniya SSSR.  
(RADIATION SICKNESS) (ALIMENTARY CANAL)

KISILEV, P.N., prof.; SEMINA, V.A.

Some immunological self-protective mechanisms of the organism  
against ionizing radiations. Vop.radiobiol. 2:356-363 '57.  
(MIRA 12:6)

1. Sotrudniki Tsentral'nogo nauchno-issledovatel'skogo rentgeno-  
radiologicheskogo instituta Ministerstva zdravookhraneniya SSSR.  
(IMMUNITY) (RADIATION--PHYSIOLOGICAL EFFECT)

KISELEV, P.M., prof.; RABINOVICH, R.M.; METER, I.D.

Course and treatment of experimental pneumonia in animals with radiation sickness. Vop.radiobiol. 2:364-372 '57.

(MIRA 12:6)

1. Sotrudniki Tsentral'nogo nauchno-issledovatel'skogo rentgeno-radiologicheskogo instituta Ministerstva zdravookhraneniya SSSR.  
(RADIATION SICKNESS) (PNEUMONIA)

*Kiselev, P.N.*

KISELEV, P.N.

Some results of studying the effect of ionizing radiations on infection and immunity [with summary in English]. Med.rad. 2 no.5:55-64 S-O '57. (MIRA 11:2)

1. Iz bakterio-serologicheskoy laboratorii (zav. - prof. P.N.Kiselev) TSentral'nogo nauchno-issledovatel'skogo rentgeno-radiologicheskogo instituta Ministerstva zdravookhraneniya SSSR.

(IMMUNITY, effect of radiations,  
ionizing, review (Rus))

(INFECTION, experimental,  
eff. of ionizing radiations, review (Rus))

(RADIATIONS, effects,  
ionizing, on exper. infect. & immun., review (Rus))

USSR/General Problems of Pathology - Immunity

U-1

Abs Jour : Ref Zhur - Biol., No. 18, 1958, 84713

Author : Kiselev, P. N.

Inst : No institute is given

Title : The Protective Role of Tissues of the Organism against its Own Denatured Proteins

Orig Pub : Zh. Mikrobiol., Epidemiol. i Immunobiol., 1957, No. 6, 104-111

Abstract : The formation of antibodies against the organism's own denatured proteins (AODP) has been demonstrated with the aid of the RSK <sup>Blood Serum</sup> <sup>Re-</sup> <sup>action</sup> in which the antigen was the animals' own serum proteins, partially denatured by three-hour heating at 60 degrees or a 24-hour exposure to the action of 50-percent alcohol. In animals with inflammation due to turpentine irritation (aseptic) or to staphylococci, the titer of AODP is higher the greater the inflammation and necrosis of the

Card 1/3

USSR/General Problems of Pathology - Immunity

U-1

APPROVED FOR RELEASE: 09/17/2001

CIA-RDP86-00513R000722810009-2"

Abs Jour : Ref Zhur - Biol., No. 18, 1958, 84713

Abstract : tissues. In a number of experiments the protective significance of the AODP was established: upon repeated administration of turpentine to the skin of guinea pigs, observation was made of the formation of AODP and a corresponding subsequent diminution in the size of the focus of inflammation and necrosis and a shortening of the healing time of the focus by an average of nine days following the second or third injection. Elevation of the resistance of mice upon repeated X-irradiation was also accompanied by the appearance of AODP in titers which were greater the higher the survival rate of the animals. Upon single irradiation of various parts of the body of mice with 130 r, there was a direct relationship between the percentage of surviving animals and the antibody titer. Immunization of the mice with homologous denatured protein led to the formation of AODP in a titer of 1:50 (on the average) and to an increase of two times in the

Card 2/2

KISELEV, P.N.; RABINOVICH, R.M.; METER, I.D.

Treatment of staphylococcal pneumonia in radiation sickness [with summary in English]. Med.rad. 3 no.4:41-46 J1-Ag '58.

(MIRA 12:3)

1. Tsentral'nogo nauchno-issledovatel'skogo rentgeno-radiologicheskogo instituta Ministerstva zdavookhraneniya SSSR.

(PNEUMONIA, experimental,

micrococcal, in radiation sickness, eff. of antibiotics (Rus))

(ANTIBIOTICS, effects,

on exper. micrococcal pneumonia in radiation sickness (Rus))

(MICROCOCCAL INFECTIONS, experimental,

pneumonia in radiation sickness, eff. of antibiotics (Rus))

(ROENTGEN RAYS, effects,

total body, eff. of antibiotics on assoc. pneumonia in animals (Rus))

KISELEV, P.N.; RABINOVICH, R.M.; METER, I.D.

Principles of the development, course and outcome of experimental pneumonia in acute radiation sickness. Med.rad. 3 no.6:3-10 N-D '58. (MIRA 12:1)

1. Iz bakterio-serologicheskoy laboratorii (zav. - prof. P.N. Kiselev) Tsentral'nogo nauchno-issledovatel'skogo rentgeno-radiologicheskogo instituta Ministerstva zdavookhraneniya SSSR.

(PNEUMONIA, exper.

eff. of exper. radiation sickness (Rus))

(ROENTGEN RAYS, effects,

on exper. pneumonia, radiation sickness dose (Rus))

KISELEV, P.N.; SIVERTSEVA, V.N.; KARPOVA, Ye. V.

~~Characteristics of the course of infectious processes as effected by~~  
ionizing irradiation of the body. Zhur. mikrobiol. epid. i imm. 29  
no.10:21-29 0 '58. (MIRA 11:12)

1. Iz Tsentral'nogo rentgeno-radiologicheskogo instituta Ministerstva  
sdravookhraneniya, SSSR.

(MICROCOCCAL INFECTIONS, exper.

~~eff. of X-rays (rus))~~

(ROENTGEN RAYS, effects,

on exper. micrococcal infect. (Rus))

KISELEV, P.N.; SEMINA, V.A.

Some immunological mechanisms of autoprotection of the organism  
from the effect of ionizing radiations. Zhur.mikrobiol.epid. i  
imun. 30 no.1:44-40 Ja '58. (MIRA 12:3)

1. Iz TSentral'nogo nauchno-issledovatel'skogo rentgeno-radiologi-  
cheskogo instituta Ministerstva zdравookhraneniya SSSR.

(IMMUNITY,

eff. of bact. toxins on radio-sensitive & radio-  
resist. animals (Rus))

(RADIATIONS, effects,  
same)

KISELEV, P.N.

**830 Experiences with Irradiation of Patients with Cerebral Tumours**  
GALPERIN, M. D. Leningrad (Soviet Union)

At present, surgical as well as irradiation treatment are regarded as being the most effective therapy. The combined methods occasionally prove to be highly beneficial.

The pre-requisite for the success of either method of treatment of tumours is early diagnosis. For the successful treatment of brain-tumours, the choice of the method is all-important. The surgical and irradiation treatments are not competitive. Taking into consideration the position of the tumour, its extent, its histological structure, the characteristics of the clinical course and the general condition of the patient, preference should be given to one or the other method.

The author examined recent and past results of irradiation and combined irradiation and surgical treatment of brain tumours during 1955-1958. 350 case-histories of patients of the Neurogen-Radiological Department of the Benkierov Institute of Neuro-Psychology were evaluated.

Irradiation was performed on patients with tumours of varied localisation and of different histological structure. Comparative assessment of the results of the different methods of irradiation in the patients was carried out, depending on the histogenesis of the tumour.

Analysis of these case-histories indicates that the elaborated and applicable methods of irradiation of brain-tumours prolong considerably the life of the patients and have an immediate, marked curative effect.

The complications occurring during treatment as well as afterwards were also studied. Clinical indications and contra-indications of irradiation of patients with brain-tumours were elaborated.

**831 Effect of the Chronic Influence of low Doses of Ionizing Irradiation on the Humoral and Cell-Linked Immunity in Animal Experiments**  
KISELEV, P. N. KISELEV, P. N. Leningrad (Soviet Union)  
BUZINT, T. A. Leningrad (Soviet Union)

The authors investigated the changes in natural immunity and immunogenic processes in laboratory animals under chronic irradiation with low doses of the gamma rays of  $^{60}\text{Co}$ . The dose-performance of the irradiation was 1.6-2.2 r/day. The period of irradiation lasted from 30 days to 2.5 years. The total dose was 50-4,500 r. The effect of these irradiation doses led to the development of chronic radiation illness. On this basis the disorder in the humoral and cellular natural immunity and immunogenicity was investigated, with the following results:

1. Under chronic, uninterrupted action, lethal radiation disease develops through auto-infection. The total lethal dose exceeds the single dose by 2-4 times. Chronic radiation illness is accompanied by negligible leucopenia, preceded by a phase of leucocytosis.
2. Under chronic irradiation natural immunity and immunogenicity are disturbed. Reduction of natural immunity appears after 9-10 months and becomes manifest by a reduction of bactericidity of the blood, leucopenia and reduction of phagocytic activity of the leucocytes. A change in the titre of the complement was observed only as late as after 10-12 months. Bacteremia is preceded by reduced bactericidity of the blood.
3. The disturbance of cellular immunity is indicated by an increased sensitivity to leukaemia, by an enhanced reproduction of virus, by a lowering of the regenerative and digestive capability of the reticulo-endothelial cells.

No. 861-882

4. The most marked reduction of natural immunity occurs in young animals born of irradiated parents and subjected during the period of embryogenesis to the effect of irradiation.

5. Chronic irradiation of an organism leads to disturbance of immunogenesis. However, as an irradiation dose equal to a single one, the production of antibodies is less suppressed. These differences are connected with the adaptation mechanisms and reparative processes in the tissues producing the antibodies.

6. The phase of suppression of natural immunity and immunogenesis may be preceded by a period of their stimulation. At a total dose of 50-100 r, the following is observed: increase of bactericidity of the blood, increased phagocytic activity of the leucocytes, of the cells of the reticulo-endothelial system; reduction of sensitivity to leukaemia; stimulation of anti-body formation.

**881 Brachytherapy of Cancer of the Oral Cavity, the Nasopharynx, and the Antrum**  
KOSLOVA, A. V. KOSLOVA, A. V. Moscow (Soviet Union)

Early results of the use of radio-active preparations (radium, radio-active cobalt, gold, chromophosphate) in the treatment of 221 patients are presented. Among these 221 patients there were 67 with malignant tumours of the oral cavity, 55 with malignant tumours of the nasopharynx and 111 patients with malignant tumours of the antrum.

Growths in the 1st and 2nd stages were found in 66 patients, stage 3 in 150 and stage 4 in 55 patients.

Treatment consisted in the combined method of ray therapy (radium surgery, cavity therapy, applicator therapy and telegamma therapy).

The patients were under observation over 3-10 years. Recovery was observed in 61% of the patients with malignant tumours in all 4 stages.

In none of the patients irradiation was followed by complication. The methods and the results of treatment are discussed.

**882 Radiological Investigations and Rational Means of Reducing the Dose**  
POBEDINSKY, M. N. Leningrad (Soviet Union)

During recent years the natural level of radiation has risen continuously. One of the factors raising this level are the radio-diagnostic examinations, which, according to reports from foreign authors, increase the amount of radiation acting on the population by 21% - 30%.

Particular attention should be given to the effect of radiation on the sex glands. Genetic sequelae of radiation may occur even with very low doses.

In X-ray exposures and fluoroscopy the focus dose of the radiation striking the sex glands may be 100-1,000 mrad. The variations in dose depend on the conditions of radiation and also on the part of the body in which radiation is directed. The highest doses affecting the sex glands occur in examinations of the pelvic region, the hip and the abdomen, especially when repeated.

The necessity for an ever-increasing extension of radio-diagnostic for the population, and the new methods of radiological examinations in practice, require research into means for reducing the radiation dose acting on the sex glands during X-ray exposures and fluoroscopy. In order to reduce the radiation dose and to prevent serious radiation effects, the following is required: highly qualified medical staff carrying out the radio-diagnostic examinations; knowledge of the radiation dose to which the patient is subjected and registration of the dose in the patients' card-index. The examinations should be performed with harder rays, using heavier filtration and increased voltage, as well as short focus, distance.

In fluoroscopy the following is required: satisfactory adaptation of the type of the examination, organization of the working hours and low unacceptance. In addition, the advantages of working with narrow beams should be made use of.

In radiological examination of the pelvic region, the hip and the abdomen, the sex glands must be protected from direct radiation.

Presented at the Ninth International Congress of Radiology, Munich, 23-30 July 1959.

KISELEV, P. N., SIVERTSEVA, V. N., KARPOVA, YE. V.

"Basic rules of development of infectious processes upon the effect of large doses of ionizing radiation on the organism."

report submitted at the 13th All-Union Congress of Hygienists, Epidemiologists and Infectionists, 1959.

KISELEV, P.N.; BUZINI, P.A.

Effect of chronic uninterrupted ionizing radiation on immunity.  
Med. rad. 4 no.4:36-44 Ap '59. (MIRA 12:7)

1. Iz bakterio-serologicheskoy laboratorii (zav. - prof. P.N. Kiselev) Tsentral'nogo nauchno-issledovatel'skogo rentgeno-radiologicheskogo instituta Ministerstva zdravookhraneniya SSSR.  
(IMMUNITY,

eff. of gamma rays on immunogenesis in animals (Rus))  
(GAMMA RAYS, effects,  
on immunogenesis in animals (Rus))

KISELEV, P.N., prof.; SMORODINTSEV, A.I. A.

"Effect of ionizing radiation on immunity" by V. L. Troitskii, M.A.  
Tumanian. Reviewed by P. N. Kiselev, A.A. Med. rad. 4 no.5:91-93  
My '59. (MIRA 12:7)  
(IMMUNITY) (RADIATION SICKNESS) (TROITSKII, V.L.)  
(TUMANIAN, M.A.)

PHASE I BOOK EXPLOITATION

SOV/5435

Kiselev, P. N., Professor, G. A. Gusterin, and A. I. Strashinin, Eds.

Voprosy radiobiologii. t. III: Sbornik trudov, posvyashchenny 60-letiyu so dnya rozhdeniya Professora M. N. Pobedinskogo (Problems in Radiation Biology. v. 3: A Collection of Works Dedicated to the Sixtieth Birthday of Professor M[ikhail] N[ikolayevich] Pobedinskiy [Doctor of Medicine]) Leningrad. Tsentr. n-issl. in-t med. radiologii M-va zdravookhrananiya SSSR, 1960. 422 p. 1,500 copies printed.

Tech. Ed.: P. S. Peleshuk.

PURPOSE: This collection of articles is intended for radiobiologists.

COVERAGE: The book contains 49 articles dealing with pathogenesis, prophylaxis, and therapy of radiation diseases. Individual articles describe investigations of the biological effects of radiation carried out by workers of the Central Scientific Research Institute for Medical Radiology of the Ministry of Public Health, USSR. [Tsentr. 'nyy nauchno-issledovatel'skiy institut meditsinskoy radiologii Ministerstva zdravookhraneniya SSSR] during 1958-59. The following

Card 1/10

Problems in Radiation Biology (Cont.)

SOV/5435

topics are covered: various aspects of primary effects of radiation; the course of some metabolic processes in animals subjected to ionizing radiation; reactions in irradiated organisms; morphologic changes in radiation disease; and reparation and regeneration of tissues injured by irradiation. Some articles give attention to the effectiveness of experimental medical treatments. No personalities are mentioned. References accompany almost all of the articles.

TABLE OF CONTENTS:

Foreword	3
Gusterin, G. A., and A. I. Strashinin. Professor Mikhail Nikolayevich Popedinskiy (Commemorating his Sixtieth Birthday)	5
Isbedinskiy, A. V. [Member, Academy of Medical Sciences USSR], N. I. Arlashchenko, and V. M. Mastryukova. On the Mechanism of Trophic Disturbances Due to Ionizing Radiation	11
Zedgenidze, G. A., [Member, Academy of Medical Sciences USSR], Ye. A. Zherbin, K. V. Ivanov, and P. R. Vaynshteyn. Hormonal Activity of the Adrenal Cortex in Acute Radiation Sickness and the Effect of Desoxycorticosterone Acetate on the Disease	17

Card 2/10

Problems in Radiation Biology (Cont.)

SOV/5435

Cherkasov, V. F. Reactions of the Cardiovascular System and the Respiration of Irradiated Animals to Some Functional Factors	35
Bryukhanov, O. A. Concerning the Problem of Depressant Reactions in Radiation Sickness	44
Lazovskaya, A. V. Effect of Strophanthin on the Heart of Irradiated Animals	50
Shitova, Z. I., and Ye. I. Komarov. On the Reflex Mechanism of the Change in Oxygen Absorption by Intestinal Tissue During Local Irradiation With Radioactive Strontium	55
Remizova, I. V. On Some Features of Functional Changes in the Nervous and Blood System During Repeated Small-Dose Irradiation	61
Traskunova, N. V. Effect of Blocking the Sympathetic Subdivision of the Vegetative Nervous System on the Development and Course of Acute Radiation Sickness	68
Card 3/10	

Problems in Radiation Biology (Cont.)

SOV/5435

Poplavskiy, K. K. Phasic Changes in the Ability of Irradiated Animals to React to Anesthetization	78
Bondarenko, I. V. On the Reaction of Irradiated Dogs to the Introduction of Alpha Dinitrophenol	86
Alekseyeva, G. N. Reaction of an Irradiated Organism to the Introduction of Gangliolytic Preparations [gangliolitiki]	93
Protas, L. R., and A. A. Danilin. The Mechanism of Functional Disturbances in the Alimentary Canal During Acute and Subacute Forms of Experimental Radiation Sickness	97
Aleksandrov, S. N. Some Methods of Approach to the Study of Early Stages of Radiation Sequelae	104
Manoylov, S. Ye. Respiration of Tissue and Sensitivity to Radiation	111
Kachur, L. A., P. N. Kiselev, and A. N. Shutko. Effect of Ionizing Radiation on the Water-Exchange Process Between the Blood and the Extravascular Liquids in the Organism	138

Card 4/10

Problems in Radiation Biology (Cont.)

SOV/5435

- Manoylov, S. Ye., and B. A. Orlov. Use of the Spectroscopic Research Method in the Study of the Condition of Iron-Containing Compounds in Animal Organisms Irradiated With X-Rays 152
- Demin, N. N., [Professor]. On Some Metabolic Disturbances in Lipides Due to External Whole-Body Ionizing Irradiation 158
- Keylina, R. Ya. Concerning the Problem of the Condition of Processes of Synthesis and Decomposition of Carbohydrates in the Animal Organism Subjected to Whole-Body X-Ray Irradiation 165
- Shitova, Z. I. Changes in Respiration of the Skeletal Muscles Following Irradiation by Radon of Different Segments of the Nervous System 173
- Mytareva, L. V. Effect of Ionizing Radiation on a Ferment of Glycolysis of Phosphohexoisomerase in Some Organs and Tissues of an Animal Organism Subjected to Whole-Body Irradiation 183

Card 5/10

Problems in Radiation Biology (Cont.)

SOV/5435

Strelin, G. S., and Ye. M. Pil'shchik. On the Repression of Regenerative Processes in the Bone Under Various Conditions of Irradiation of Animals	188
Pushnitsina, A. D. Reactive Changes in Rat Marrow in Radiation Sickness Complicated by Loss of Blood	203
Gankina, K. I., A. I. Strashinin, G. S. Strelin, and I. V. Shiffer. Regeneration of Surgical Intestinal Wounds in Radiation Sickness	211
Shmeleva, N. I. Effect of Ether Anesthesia on the Regeneration Process of Hematogenesis in Rats With Combined Injuries	222
Lyalin, Ye. A. Change in Some Aspects of the Activity of the Thyroid Glands Following Whole-Body Irradiation	231
Lyalin, Ye. A. Effect of Whole-Body Irradiation on the Proliferating Capability of Thyroid Tissue in White Mice	236
Kantin, A. V., and P. V. Sipovskiy. Formation of the Surgical Stump of Limb During Experimental Radiation Disease in Rabbits	241

Card 6/10

Problems in Radiation Biology (Cont.)

SGV/5435

Sipovskiy, P. V., and A. V. Kantin. Morphologic Healing Characteristics of the Amputational Bone Stump of Rabbits During Radiation Sickness and After Recovery	248
Funshteyn, L. V. On Morphogenesis in Experimental Acute Radiation Sickness	256
Fedorov, Yu. A. Effect of Fractional Whole-Body X-Ray Irradiation of Hard Dental Tissue in White Rats	265
Funshteyn, L. V., and E. I. Shcherban'. On the Phagocytic Capability of Some Segments of the Reticuloendothelial System Following Whole-Body X-Ray Irradiation of White Mice	272
Ochinskaya, G. V., and L. V. Funshteyn. Morphologic Changes in Internal Organs of Pregnant Animals Subjected to Whole-Body Irradiation	281
Kholin, V. V. Comparative Data on Injury to the Spleen in Acute Radiation Disease, Depending on Age of the Animals	288

Card 7/10

Problems in Radiation Biology (Cont.)

80W/5435

- Kashchenko, L. A., N. K. Shmidt, and P. I. Ostrovskaya-Zakharevich. Reaction of the Spleen, Mucous Intestinal Membrane, and Testicles of Frogs to the Effect of Ionizing Radiation in Whole-Body and Local Irradiation 298
- Kashchenko, L. A., P. I. Ostrovskaya-Zakharevich, and N. K. Shmidt. Reparation of Radiation Injury in Frog Testicles 311
- Kalashnikov, B. P., and Yu. S. Kaminskaya. Experimental Data on the Injurious Effect of X-Rays on the Retina Due to Local and Whole-Body Irradiation 318
- Kiselev, P. N., and V. A. Semina. Effect of Some Hormones of the Adrenal and Pituitary Glands on the Course of Autoinfectious Processes in Radiation Sickness 327
- Sivertseva, V. N. Problem of the Effect of Chronic Continuous Influence of Ionizing Radiation on the Course of Infectious Processes 335
- Smorodintsev, A. A. Morphologic Changes in the Respiratory Canal in Experimental Influenza of Immune White Mice Irradiated With X-Rays 344

Card 8/10

Problems in Radiation Biology (Cont.)

SOV/5435

- Kashkin, K. P. On the Possibility of Adaptation of Bacterium Faecalis  
Alcaligenes to the Effect of Ionizing Radiation 350
- Meter, I. D. Some Data on Causes of Unsuccessful Treatment of  
Radiation Disease With Antibiotics 360
- Rabinovich, R. M. X-Ray-and-Anato. \ Characteristics of Pulmonary  
Changes in Experimental Staphylococ. c Pneumonia of Irradiated Animals 369
- Petrov, I. R. [Member, Academy of Medical Sciences USSR], V. A. Bondina,  
and I. V. Il'inskaya. Use of the Dextran-Type Synthetic Colloidal  
Solution in Combined Therapy of Radiation Sickness 376
- Rusanov, A. M., G. A. Bol'shakova, and V. D. Lyashenko. Effect of  
Gangliolytic Preparations [gangliolitiki] on the Course and Outcome  
of Experimental Radiation Sickness 386

Card 9/10

Problems in Radiation Biology (Cont.)

SOV/5435

Poplavskiy, K. K. Prophylactic and Therapeutic Effect of  
Anesthetic Agents in Radiation Sickness

394

Ivanova, N. A. Comparative Estimate of Protective Properties  
of Some Mercapto Compounds

406

Shcherban', E. I. Effect of Moderate Physical Exertion During  
X-Ray Irradiation on the Course of Acute Radiation Sickness

412

AVAILABLE: Library of Congress (QB652.V6)

Card 10/10

AC/rn/mas  
10-27-61

KISELEV, P.N.; NAKHIL'NITSKAYA, Z.N.

Some results of a study of the effect of ionizing radiations on  
tissue permeability. Med. rad. 5 no.9:73-82 S '60. (MIRA 13:12)  
(TISSUES—PERMEABILITY) (RADIATION—PHYSIOLOGICAL EFFECT)

KISELEV, P.N.; KARPOVA, Ye.V.; SIVERTSEVA, M.N.

Disorders of the humoral mechanism in detoxication of the organism  
in ionizing radiation injuries. Med. rad. 5 no.11:30-36 N '60.

(MIRA 13:12)

(RADIATION SICKNESS)

(TOXINS AND ANTITOXINS)

KISELEV, P.N.; SIVERTSEVA, V.N.; NIKITINA, K.I.

Detoxication disorders of the body in radiation sickness. Med.  
rad. 6 no.4:41-48 '61. (MIRA 14:12)  
(RADIATION SICKNESS) (TOXINS AND ANTITOXINS)

KISELEV, P.N.; KASHKIN, K.P.; BOLTAKS, Yu.B.; VITOVSKAYA, G.A.

Acquisition of resistance to radioactivity by a microbe cell kept  
in a medium with a high natural radiation level. Mikrobiologiya  
30 no.2:207-213 Mr-Apr '61. (MIRA 14:6)

1. Tsentral'nyy nauchno-issledovatel'skiy institut meditsinskoy  
radiologii i Khimiko-farmatsevticheskiiy institut, Leningrad.  
(BACTERIA) (RADIATION—PHYSIOLOGICAL EFFECT)

KLSA ELEV, P. N.

(6)  
Changes in Cellular Immunity Under the Influence of Chronic Continuous Exposure to Relatively Small Doses of Ionizing Radiation 2

P. N. Klychev and P. A. Boyin

The influence of chronic continuous exposure to small doses of  $^{60}\text{Co}$   $\gamma$ -rays on the humoral and cellular immunity of the organism has been studied. The investigations were carried out on different experimental animals (mice, rats, guinea-pigs and rabbits). The dose rate varied from 0.5 to 3.5 r/day. Irradiation was protracted from 30 days to 5 yr. The total dose was 100-2000 r.

Such an exposure resulted in the development of chronic radiation disease manifested in different degrees. The observed changes in the humoral immunity were of a phasic type and manifested themselves in stimulation and subsequent inhibition which were accompanied by the development of auto-infectious processes.

The experiments have shown that chronic continuous exposure to small doses of ionizing radiation brings about a disturbance of the cellular immunity. These changes also are of a phasic type and characterized by the same regularities. The increase in the immunity is always less manifest than its subsequent inhibition. The decrease in the cellular immunity is expressed: (1) in reduction of function of the reticulo-endothelial system, occurring rather late also in the case of sufficiently large total doses; (2) in changes of absorbing and digesting functions of the leucocytes; (3) in increasing the sensitivity of the organism towards exo- and endo-toxins; and (4) in changing the capability of somatic cells to remove heterogeneous particles and to inhibit the growth of viruses. A certain stimulation of the organism preceding its weakening is revealed in a lesser delay of the growth of influenza virus in the organism, stimulation of a RES function and increasing the phagocytic capability of the leucocytes.

The Central Research Institute of Medical Radiology of the Ministry of Public Health of the USSR, Leningrad.

100

report presented at the 2nd Intl. Congress of Radiation Research,  
Harrogate/Yorkshire, Gt. Brit. 5-11 Aug 1962

KISELEV, P.N.

Influence of the endocrine system on the localization of infectious processes in the body. Trudy Len.khim.-farm.inst. no.13:57-62 '62.

(MIRA 15:10)

1. Kafedra mikrobiologii (zav. prof. P.N.Kiselev) Leningradskogo khimiko-farmatsevticheskogo instituta.

(ENDOCRINE GLANDS) (INFECTION)

40924

27.1220

S/241/62/007/008/001/001

I015/I215

AUTHOR: Kisilev, P. N. and Buzini, P. A.

TITLE: Changes in cell immunity upon chronic continuous exposure to ionizing radiation

PERIODICAL: Meditsinskaya radiologiya, v. 7, no. 8, 1962, 59-65

TEXT: The effect of chronic ionizing radiation on the mechanism of immunity has not been sufficiently studied. This is the continuation of a previous study. Rabbits, guinea pigs, albino mice, and rats were subjected to an irradiation of 50-4000 r with Co<sup>60</sup> gamma rays at 0.5-4.3 r/24 hrs. The observations continued for 5 years. A dose of 465 r did not alter the properties of WBC, RES, and somatic cells. The phagocytosing and digesting properties of WBC were distorted after larger doses of irradiation and up to a dose of 700-1000 r there was a definite dependence between these factors. Irradiation during embryogenesis brought about the development of nonphagocytosing WBC, even after a total dose of 200 r. The RES function was studied with S<sup>35</sup>-labelled B. coli and colloidal Au<sup>198</sup>. The RES function was evaluated by the method of Benacerraf, Halpern, et al. A chronic continuous irradiation at relatively small doses (2.4 r/24 hours) brings about a suppression of the RES function following a prior stimulation. The sensitivity of somatic cells to bacterial toxins and viruses was also altered in chronic continuous irradiation (950 r at 1.29 r/24 hrs). Sensitivity was estimated by determination of LD<sup>50</sup> in both control and experimental animals. The increased sensitivity

Card 1/2

Changes in cell immunity ...

S/241/62/007/008/001/001  
I015/I215

of cells to endo- and exotoxins was preceded by a decreased sensitivity (up to a total dose of 285 r). Thus, chronic irradiation with small doses affects both the humoral and cellular immunity simultaneously and with the same characteristics. There are 2 figures and 4 tables.

**ASSOCIATION:** Laboratoriya radiatsionnoy mikrobiologii i immunologii (zav.-prof. P. N. Kiselev) Tsentral'nogo nauchno-issledovatel'skogo instituta meditsinskoy radiologii (dir.-kandidat meditsinskikh nauk E. I. Vorob'yev), Ministerstva zdravookhraneniya SSSR (Laboratory of Radiation Microbiology and Immunology [headed by Prof. P. N. Kiselev], Central Scientific Research Institute of Medical Radiology [directed by Candidate Medical Sciences E. I. Vorob'yev], Ministry of Health, USSR.)

**SUBMITTED:** February 14, 1962

Card 2/2

L 13066-63

EWA(b)/EWT(1)/EWT(m)/BDS AMD/ASD/AFFTC/APGC Pa-4 AR/K

ACCESSION NR: AP3000257

S/0241/63/008/005/0033/0039

AUTHOR: Kiselev, P. N. (Director); Sivertseva, V. N.

TITLE: Change in capacity of irradiated animal tissues to destroy  
microbe toxins

SOURCE: Meditsinskaya radiologiya, no. 5, 1963, 33-39

TOPIC TAGS: radiation sickness, microbe toxin, endotoxin, spleen,  
reticuloendothelial system, detoxicating mechanism

ABSTRACT: An organism with radiation injuries is highly sensitive to microbe toxins, particularly to endotoxins in tissues, because its detoxication mechanisms are impaired. Humoral factors are important in exotoxin destruction in the blood, but how can the endotoxins in tissues be destroyed? Sources in the literature have suggested that the reticuloendothelial system (15% of the body) may play an important role. This is a study of the effect of the reticuloendothelial system as represented by mice spleens on endotoxin destruction in tissues. Mice were subjected to sublethal doses of X-ray irradiation and spleen extracts were prepared. Dry endotoxin (S. Breslav type) was dissolved in extract solutions and incubated. Cortisone in

Card 1/2

L 13066-63

ACCESSION NR: AP3000257

varying doses was used to stimulate the spleen. Results show that the spleen has a remarkable capacity for destroying endotoxins in pure form or in the form of microbe bodies. Spleen extract prepared from mice at the height of radiation sickness is two times less effective in destroying endotoxins than extracts from nonirradiated mice. Use of cortisone strengthens the detoxicating mechanisms and makes them more resistant to endotoxins (Figs. 4, 5, 6). The author concludes that the reticuloendothelial system's capacity to destroy endotoxins is apparently of an enzyme nature. Endotoxins can be destroyed in the tissues by activating natural detoxication mechanisms with cortisone (and probably other stimuli), but only to a certain degree because in advanced stages of radiation sickness the mechanisms are totally impaired. Orig. art. has: 6 figures.

ASSOCIATION: Laboratoriya radiatsionnoy mikrobiologii i immunologii, Tsentral'nogo nauchno-issledovatel'skogo instituta meditsinskoy radiologii, Ministerstva zdravookhraneniya SSSR (Laboratory of Radiation Microbiology and Immunology of the Central Scientific Research Institute of Nuclear Medicine of the Ministry of Public Health SSSR)

Card 2/32

ACCESSION NR: AP4038943

8/0241/64/000/005/0050/0058

AUTHOR: Kiselev, P. N.; Buzini, P. A.

TITLE: Remote immunologic sequelae of the effect of ionizing radiation

SOURCE: Meditsinskaya radiologiya, no. 5, 1964, 50-58

TOPIC TAGS: immunologic sequelae, ionizing radiation, immunogenesis, immunity retention, Salmonella enteridis Gartneri, Salmonella breslau, Pseudomonas aeruginosa, endotoxin sensitivity, bacteremia, phagocytosis, delayed postirradiation infection, mononuclear phagocyte, polynuclear phagocyte, infection resistance, antibody synthesis, vaccination

ABSTRACT: The duration of retention of natural immunity and immunogenesis acquired during acute radiation sickness was studied in guinea pigs and white mice which received 200 r (LD<sub>30/30</sub> days) and 465-500 r (LD<sub>50/30</sub>) respectively. Resistance to infection was tested 3-28 months after irradiation, in either spontaneously infected animals or by infection with Salmonella enteridis Gartneri, Salmonella breslau or Pseudomonas aeruginosa. Endotoxin sensitivity was investigated by introducing

Card | 1/4

ACCESSION NR: AP4038943

the endotoxin in a LD<sub>50/3</sub> days dose and the bacteremia determined by microbial count in blood and organs. The number of leucocytes in peripheral blood, phagocytic activity in peritoneal exudate, etc. served as immunologic indicators. Spontaneous infection of the mice with *Salmonella enteridis* Gartneri after 5½ months led to the death of 58% irradiated against 18% of non-irradiated animals. Experimental infection with 100 million bodies of the same agent in 380 mice, half of which had been irradiated 6 months earlier, led to death of 72% of the irradiated, 38% of the control mice. *Salmonella breslau* infection under the same conditions yielded the corresponding figures of 97 against 65%. Spontaneous infection of guinea pigs with *Pseudomonas aeruginosa* 19 months after irradiation led to death of 32 against 14%. Experimental infection with the same agent in a group of mixed age was performed 3, 6, 9, 12, 19 and 28 months after irradiation. Increased sensitivity to infection (compared to controls) decreased only after the 19th month, to return to normal after 28 months. A study of the reasons for such increased sensitivity comprised testing the bactericidal property of the blood against *Salmonella typhi* murium, the complement titer, phagocytic activity of leucocytes, bacteremia in mice with acute and chronic radiation sickness, as well as in controls 6 months after irradiation. Microbe lysing activity was found decreased

Card 2/4

ACCESSION NR: AP4038943

despite close to normal counts. Similar studies with *Pseudomonas aeruginosa* in guinea pigs 3-28 months after irradiation showed e.g. a decrease in mononuclear and increase in polynuclear phagocytes, their numbers returning to normal only after 28 months. Microbial count in guinea pig blood at various periods following irradiation confirmed the decreased phagocytic activity as late as 19 months after irradiation. Mouse sensitivity to *Salmonella typhi* abdominalis endotoxin was 2.6 times that of controls after 2½ months, 1.6 times after 5½ months. The development of antibodies to *Salmonella breslau* under these conditions was determined in guinea pigs vaccinated 3, 12 and 19 months after irradiation. No disturbance of immunogenesis was found; thus antibody synthesis is only disturbed during acute radiation sickness. Decreased anti-infective properties of the tissues however, although not irreversible, persist during half of the life span of guinea pigs and white mice. Orig. art. has: 5 tables and 6 figures.

ASSOCIATION: Allergologicheskaya laboratoriya Nauchno-issledovatel'skogo instituta ukha, gorla i nosa (Allergy Laboratory, Scientific Research Institute of Ear, Throat and Nose)

Card | 3/4

KISELEV, P.N.; NIKITINA, K.I.; CHEN SHAO-CHAN; KARMANOVA, Z.P.

Role of autoinfection in the development of hemorrhagic syndrome  
in acute radiation sickness. Radiobiologiya 4 no.5:790-793 '64.

(MIRA 18:4)

1. Tsentral'nyy nauchno-issledovatel'skiy rentgeno-radiologicheskiy  
institut Ministerstva zdravookhraneniya SSSR, Leningrad.

KISELEV, P.N.; SEMINA, V.A.

Ways of normalizing immunogenesis disturbed by the action of ionizing radiations. Med.rad. 9 no.9:61-67 S '64.

(MIRA 18:4)

1. Laboratoriya radiatsionnoy immunologii (zav. - prof. P.N. Kiselev) Tsentral'nogo nauchno-issledovatel'skogo rentgenoradiologicheskogo instituta (dir. Ye.I.Vorob'yev) Ministerstva zdavookhraneniya SSSR.

KISELEV, P.N.; NIKITINA, K.I.; CHEN SHAO-CHAN

Significance of the formation of antiendotoxins against *Escherichia coli* in the involution of hemorrhagic syndrome in radiation sickness.  
Radiobiologiya 5 no.1:87-92 '65. (MIRA 18:3)

1. Tsentral'nyy nauchno-issledovatel'skiy institut meditsinskoy radiologii Ministerstva zdravookhraneniya SSSR, Moskva.

KISELEV, P.N.; KARPOVA, Ye.V.

Role of the sulfhydryl (SH) groups of proteins in the block and the fixation of the complement. Zhur. mikrobiol., epid. i immun. 41 no.11:43-48 '65. (MIRA 18:5)

1. TSentral'nyy nauchno-issledovatel'skiy institut meditsinskoy radiologii Ministerstva zdravookhraneniya SSSR.

KISELEV, P.N.; KARPOVA, Ye.V.

Significance of the changes in the activity of tissue hyaluronidase in disorders of tissue permeability under the effect of ionizing radiation. Med. rad. 10 no.1:54-61 Ja '65. (MIRA 18:7)

1. Tsentral'nyy nauchno-issledovatel'skiy institut meditsinskoy radiologii Ministerstva zdravookhraneniya SSSR, Leningrad.

I 0057-67 EWT(1) JK

ACC NR: AP6033869

SOURCE CODE: UR/0205/66/006/005/0763/0765

AUTHOR: Kiselev, P. N.; Sivertseva, V. N.

ORG: X-Ray Radiological Institute, MZ SSSR, Leningrad (Rentgeno-radiologicheskiy institut MZ SSSR)

TITLE: Effects of oxidation and phosphorylation on the microbial toxin sensitivity of irradiated animals

SOURCE: Radiobiologiya, v. 6, no. 5, 1966, 763-765

TOPIC TAGS: biologic oxidation, biologic phosphorylation, metabolic effect, irradiation, toxin, microbial toxin, toxin effect, *toxicology*, *radiation biochemical effect*

ABSTRACT: Disturbance of oxidation and phosphorylation processes is one of the possible reasons for the increased sensitivity of an irradiated animal to toxins. Reducing substances also lower toxin resistance. Animals receiving injections of alpha-dinitrophenol became more resistant to endotoxin. Orig. art. has: 1 table. [W.A. 50]

SUB CODE: 06/ SUBM DATE: 09Mar65/ ORIG REF: 008/ OTH REF: 003

UDC: 577.391:612.017.1

Card 1/1

KISELEV, P.P.

86-9-14/36

AUTHOR: Kiselev, P.P., Col.

TITLE: Bombing the Small-Size Targets from a Fighter-Bomber  
(Bombometaniye s istrebitelya-bombardirovshchika po  
malorazmernym tselyam)

PERIODICAL: Vestnik Vozdushnogo Flota, 1957, Nr 9, pp. 40-46 (USSR)

ABSTRACT: In this article the author recommends that a sighting-  
aiming device should be installed in fighter-bombers,  
in order to train the flying personnel in accurate bomb-  
ing of point targets and small-size targets. The bombing  
from medium and low altitudes at diving angles of 30 - 60°  
and at lead angles of 5 - 11° is difficult, because the  
point of aim, when ASP-3n sight is used, should be moved  
forward from the center of the target by the amount of  
lead allowance. In view of the fact that the aiming  
allowance changes within a wide range (from 200 - 800 m),  
when point targets or targets of small dimensions are  
bombed, the visual determination of the aiming point is  
associated with considerable errors. The device, suggested  
by the author, permits to aim directly at the center of  
the target regardless of its size. The device consists of

Card 1/2

APPROVED FOR RELEASE: 09/17/2001 CIA-RDP86-00513R000722810009-2

Bombing the Small-Size Targets from a Fighter-Bomber (cont.)

the sighting frame installed in the pilot's cabin, of a  
pin for the directional control which is fastened to the  
upper cowl of the nose part of the fuselage, and of  
a vertical course line with marks on the front bullet-  
proof glass of the cabin enclosure. Among others, it is  
mentioned that the bomb fuzes, type MDV-1, operate with-  
out fail when the time of bomb fall is not lower than  
11 - 11.5 seconds. Four diagrams, 1 table.

AVAILABLE: Library of Congress.

Card 2/2

KISELEV, P.P.

Specifications for part drawings. Standartizatsia 27 no.10:  
46 0 '63. (MIRA 16:11)

KISELEV, P.S., ~~pyardii~~ general-major aviatsii

Birth of a guards regiment. Vest. Vozd. Fl. 41 no. 6:66-73 Je '58.  
(MIRA 11:7)

(Aeronautics, Military)  
(Fighter planes)

KISELEV, P.V.

Our workers are our gold assets. Avtom., telem. i sviaz' 8  
no.4:19-21 Ap '64. (MIRA 18:2)

1. Nachal'nik Bureyskoy distantssii signalizatsii i svyazi  
Zabaykal'skoy dorogi.

KISELEV, P.Ya. (Kiyev)

Approximation of analytic functions by polynomials in a finite  
number of regions. Ukr.mat.zhur. 14 no.2:202-205 '62.

(MIRA 15:11)

(Functions, Analytic) (Polynomials)

KISELEV, ~~413~~ P. Ya

S/021/63/000/001/002/012  
D251/D308

AUTHOR: Kysel'ov, P. Ya.

TITLE: On the approximation of analytic functions in the mean in a finite number of regions

PERIODICAL: Akademiya nauk Ukrayins'koyi RSR. Dopovidi, no. 1, 1963, 6-9

TEXT: The author considers a region  $D$  of the complex  $z$ -plane, bounded by a closed rectifiable Jordan curve  $\Gamma$ . If a function  $f(z)$  is analytic at interior points of  $D$  and possesses in  $D$  a  $k$ -th derivative  $f^{(k)}(z)$  belonging to class  $E_p$  in  $D$  and satisfying the generalized Lipschitz condition of order  $\alpha$  on  $\Gamma$ , the function is said to be of class  $E(k, p, \alpha)$ . A set  $E$  bounded by a finite number of mutually external smooth Jordan curves  $\Gamma_j$ ,  $j = 1, 2, \dots, \nu$  on each curve of which the angle  $\theta(s)$  which the tangent makes with the real axis has, considered as a function of the arc-length  $s$ , a modulus

Card 1/3

On the approximation of ...

S/021/63/000/001/002/012  
D251/D308

of continuity  $j(h)$  which satisfies

$$\int_0^c \frac{j(h)}{h} dh < \infty \quad (1)$$

It is proved that if a function  $f(z)$  is of class  $E(k, p, \alpha)$ ,  $p > 1$ ,  $0 < \alpha \leq 1$  in such as set  $E$ , then there exist polynomials  $\Pi_n(z)$  of degree  $n$  which satisfy

$$\|f(z) - \Pi_n(z)\|_F = \left\{ \int_F |f(z) - \Pi_n(z)|^p |dz| \right\}^{\frac{1}{p}} < \frac{M_3}{n^{k+\alpha}}$$

where  $M_3$  is a constant which is dependent only on  $E$  and  $p$ . The proof is based on the use of a function  $w = \varphi_j(z)$  which effects a

Card 2/3

On the approximation of ...

S/021/63/000/001/002/012  
D251/D308

one-one conformal transformation of the outside of  $\Gamma_j$  onto the outside of the unit circle, on Walsh's lemma, and on Mynkivs'kyi's inequality.

ASSOCIATION: Instytut matematyki AN URSR (Institute of Mathematics of the AS UkrSSR)

PRESENTED: by Yu. O. Mytropol's'kyi, Academician

SUBMITTED: September 12, 1962

Card 3/3

KISELEV, P.Ya.

Some problems involving harmonic interpolation. Trudy Sem. po  
pr.kl. mat. 1 no.1:72-83 '63. (MIRA 18:2)

1. Institut matematiki AN UkrSSR, Kiyev.

KISELEV, P.Ya. (Kiyev)

Approximation of analytic functions by Faber-Walsh polynomials.  
Ukr. mat. zhur. 15 no.2:193-199 '63. (MIRA 16:9)

KISELEV, R.A.; NIKOLAYEV, V.I.

Electromyographic picture of the functional transformation of the motor apparatus following tendon transplantation. Trudy Inst. fiziol. 6:75-85 '57. (MIRA 11:4)

1. Khirurgicheskiy sektor (zaveduyushchiy V.I. Sazonov), Laboratoriya elektrofiziologii (zaveduyushchiy V.Ye. Delov) i Travmatologicheskii institut im. R.R. Vredens (direktor V.I. Sazonov). (MUSCLES--TRANSPLANTATION)

LEVI, M.I.; GUSEV, V.M.; KISLYAKOVA, L.N.; CHUYEVA, G.I.; KISELY, R.I.; DERKACH, V.S., professor, ispolnyayushchiy obyazannost' direktora; ABRAMOV, S.G., zavednyushchiy.

Natural nidi of lymphocytic choriomeningitis. Zhur.mikrobiol.epid.i immu.  
no.8:76-81 Ag '53. (MLRA 6:11)

1. Khar'kovskiy institut epidemiologii im. I.I.Mechnikova (for Derkach).
2. Khar'kovskaya protivochumnaya stantsiya (for Abramov).  
(Meningitis, Cerebrospinal)

KISELEV, R.I.

Certain problems in the epidemiology of vesicular (pox-like)  
rickettsiosis. Zhur.mikrobiol.epid.i immun. no.1:46 Ja '54.  
(MLRA 7:2)

1. Iz Khar'kovskogo instituta epidemiologii i mikrobiologii im.  
Mechnikova.  
(Rickettsia)

KISHINEV, R.I.; VOLCHANETSKAYA, G.I.

Overall eradication of rodents and insects as a method of control of rickettsialpox. Zhur. mikrobiol. epid. i immun. no.12:28-33 D '54.

(MIRA 8:2)

1. Is Khar'kovskogo instituta vaktsin i syvorotok imeni Mechnikova  
(dir. kandidat miologicheskikh nauk B.P.Cherkas)

(RICKETTSIALPOX, prevention and control,  
continuous disinfect. & deratization)

(RATS,  
control in rickettsialpox prev.)

VOLCHANETSKAYA, G.I.; KISHLEV, R.I.

Biology of *Allodermanyssus sanguineus* Hirst. Zool.zhur. 34 no.5:  
1090-1093 S-O '55. (MLRA 9:1)

I.Khar'kovskiy nauchno-issledovatel'skiy institut vaktsin i  
syvorotok imeni Mechnikova.  
(Mites)

KISELEV, R. I., Doc of Med Sci -- (diss) "Small-pox-like rickettsiosis (Etiology, epidemiology, and measures for control)." Khar'kov, 1957, 114 pp (Khar'kov Scientific Research Institute of Vaccine and Sera im Mechnikov), 200 copies (KL, 35-57, 108)

KISELEV, R.I., BREKKE, M.L.

An allergic component in Rickettsia causing a pox-like rickettsial infection [with summary in English]. Vop.virus. 3 no.3:142-145  
My-Je '58 (MIRA 11:7)

1. Khar'kovskiy nauchno-issledovatel'skiy institut vaktsin i syvorotok imeni Mechnikova.

(RICKETTSIA,

allergic fraction causing pox-like rickettsial infect. on inoculation into guinea pigs (Rus))

LESHCHENKO, P.D., kand.med. nauk, otv. red.; CHERKAS, G.P., prof.,  
red.; PALANT, B.L., prof., red.; PEDENKO, A.I., kand.  
med. nauk, red.; KISELEV, R.I., doktor med. nauk, red.;  
KOSHEL', N.G., red.

[Diphtheria; transactions] Difteriia; sbornik trudov. Kiev,  
Gosmedizdat USSR, 1963. 155 p. (MIRA 17:6)

1. Respublikanskaya nauchno-prakticheskaya konferentsiya po  
likvidatsii difterii v USSR. 2. Ministerstvo zdravookhrane-  
niya Ukr.SSR (for Leshchenko). 3. Khar'kovskiy nauchno-  
issledovatel'skiy institut vaktsin i syvorotok im. I.I.  
Mechnikova (for Pedenko).

KISELEV, Rostislav Il'ich; ZATULOVSKIY, B.G., red.

[Rickettsialpox] Osgopodobnyi rikketsioz. Kiev,  
Zdorov'e, 1964. 103 p. (MIRA 17:11)

KHAYKINA, A.S.; DUBRAVINA, G.I.; RACHINSKAYA, A.Z.; PETRENKO, M.D.; MITEL'MAN, P.M.; KHODOROVA, Z.N.; KATS, F.M.; KISELEV, R.I.; GAYDAMAKA, M.G.; VOLOVICH, B.I.; BEKKER, M.L.; GORDIYENKO, Ye.G.; VYSOCHINENKO, Ye.K.; TELESHEVSKAYA, M.A.; NAYDEROVA, Yu.T.

Production of the active fraction of hyperimmune horse sera by means of the alcohol precipitation method under a low temperature. Nauch. osn. proizv. bakt. prep. 10:159-167 '61. (MIRA 18:7)

1. Khar'kovskiy institut vaktsin i syvorotok im. Mechnikova.

KISELEV, R.M.

All-metal saw-toothed card clothing with decreased height of teeth on the main drum of the carding machine. Izv.vys.ucheb.zav.; tekhn. tekst.prom. no.2:75-79 '58. (MIRA 11:5)

1. Ivanovskiy tekstil'nyy institut.  
(Carding machines)

KISELEV, R.M.

Saw-toothed mounting with lowered tooth height used on receiving  
drums of carding machines. Izv. vys. ucheb. zav.; tekhn. tekst.  
prom. no. 3:82-87 '58. (MIRA 11:7)

1. Ivanovskiy tekstil'nyy institut.  
(Carding machines)

KISELEV, R.M., inzh.

Operating conditions of the winding cylinder on a flat carding  
machine. Tekst.prom. 18 no.4:21-23 Ap '58. (MIRA 11:4)  
(Carding machines)

KISELEV, R.M.

Studying the expediency of the installation of two doffer  
comb pairs under the taking-in roll of a small carding  
machine. Izv. vys. ucheb. zav.; tekhn. tekst. prom. no.4:  
68-74 '63. (MIRA 16:11)

1. Ivanovskiy tekstil'nyy institut imeni M.V. Frunze.

KISELEV, R.M.

More about the design of the fillet for the receiving drums  
of carders. Izv. vys. ucheb. zav.; tekhn. tekst. prom.  
no.1:60-65 '64. (MIRA 17:5)

1. Ivanovskiy tekstil'nyy institut imeni Frunze.

SOV/85-58-9-22/33

**AUTHOR:** Kiselev, S., Sportsman 1st Rank (Sverdlovsk)

**TITLE:** New Sports Parachute (Novyy sportivnyy parashyut)

**PERIODICAL:** Kryl'ya rodiny, 1958, Nr 9, pp 24-25 (USSR)

**ABSTRACT:** The author discusses the essential features in a sports parachute and refers to the general opinion that at high altitudes a parachute requires an easily managed canopy to provide adequate speed in horizontal shifts. At low altitudes, the canopy of such a parachute should be able to develop greater or lesser speed when shifting in any direction at the will of the parachutist. It is also generally agreed that the T-2 parachute is well adapted for maneuvering and approaching a target from a rela-

Card 1/2

SOV/85-58-9-22/33

**New Sports Parachute**

tively high altitude, while the PT-1 parachute has greater target accuracy at low altitudes. The author describes his own improved version of the PT-1 parachute cupola as combining both features. There are 6 drawings.

Card 2/2

S/085/60/000/009/003/003  
A153/A029AUTHOR: Kiselev, S., Master of SportsTITLE: A New Sport Parachute 2PERIODICAL: Kryl'ya rodiny, 1960, No. 9, pp. 20-21

TEXT: 28 Having enumerated a number of shortcomings of the standard Soviet T-2 (T-2) sport parachute, the author praises the advantages of a double-slot-  
ted PT-1-2 sport parachute: a greater speed of horizontal movement and angular rotation, an elimination of sidewise "drift" of the canopy during turns, a higher suitability for jumps aiming at accuracy of landing at the target, etc. The canopy of this novel parachute is round, consisting of 28 conical panels, each one sewed together from four sections. It has a round central vent, reinforced upper and lower brims, 14 shroud lines running through the whole canopy in radial seams. The parachute has two T-shaped slots located between the 3d-5th and 24th-26th shroud lines (Fig. 1). The control shroud lines are 1,300 mm long, attached to shroud lines No. 5 and 24. By using the front harness straps it is possible to quickly turn about (to the left by using the right strap and vice versa). The use of the rear straps is not effective unless they are braced by

Card 1/3

A New Sport Parachute

S/085/60/000/009/003/003  
A153/A029


two 2m-long additional shroud lines: one such line connects the shroud line No. 26 with the semi-ring of the right rear strap, the other connects the shroud line No. 3 with the semi-ring of the left rear strap. The pilot parachute is round; the parachute pack is made of a caprone cloth. Tested by a state commission, the novel parachute has shown the following characteristics: speed of descent 5.2m/sec., speed of horizontal movement in the air 3.5m/sec., one turn round the vertical axis 5-7 sec. The sensitivity of this parachute calls upon its user to be very attentive and capable of quickly evaluating his position with respect to the ground. There are 3 figures. ✓

Card 2/3

A New Sport Parachute

S/085/60/000/009/003/003  
A153/A029

Figure 1:



Card 3/3

KISELEV, S., master sporta

The stratosphere submits to the brave. Kryl. rod. 16 no.12:8-10  
D '65. (MIRA 18:12)

1535/56 S.I.

BARKHATOV, G.V.; VASIL'YEV, V.G.; KISEL'EV, S.I.; TIKHOMIROV, Yu.P.

Oil- and gas-bearing potential of the Verkhoyansk piedmont fault  
and basic trends in prospecting this region. Geol. nefti i no. 4:  
1-7 Ap '57. (MIRA 10:8)

(Verkhoyansk Range--Petroleum geology)  
(Verkhoyansk Range--Gas, Natural--Geology)

BARKHATOV, G.V.; VASIL'YEV, V.G.; GRISHIN, G.L.; KARASEV, I.P.; KISELEV,  
S.I.; KRAVCHENKO, Ye.V.; MORDOVSKIY, V.T.; TIKHOMIROV, YU.P.;  
CHEPIKOV, K.R.; YUNGANS, S.M., ved.red.; FEDOTOVA, I.G., tekhn.red.

[Oil and gas in the eastern Siberian Platform] Neftegazonosnost'  
Vostochno-Sibirskoi platvornyy. Pod red. K.R. Chepikova. Moskva,  
Gos.nauchno-tekhn.izd-vo neft. i gorno-toplivnoi lit-ry, 1958.  
130 p. (MIRA 12:1)

1. Chlen-korrespondent AN SSSR (for Chepikov).  
(Siberian Platform--Gas, Natural)  
(Siberian Platform--Petroleum)

KRASTOSHEVSKIY, L.S., inzh.; SHEVTSOV, A.Yu., inzh.; KISELEV, S.I., inzh.

Searching for a design of precast reinforced concrete lining  
for shafts sunk by boring. Trudy VNIIONSHSa no.15:64-93 '64.  
(MIRA 18:2)

KISELEV, S. K.

Oborudovanie i ekspluatatsiia rudnichnykh pod"emnykh ustanovok. Moskva, Gostoptekhizdat, 1943, 239 p. illus.

Bibliography: p. 237.

Equipment and operation of mine hoisting units.

DLC: TN339.K5

SO: Manufacturing and Mechanical Engineering in the Soviet Union, Library of Congress, 1953.

KISELEV, S.M.

For an increase in labor productivity. Transp. stroi. 8 no.11:7  
N '58. (MIRA 12:1)  
(Machine-shop practice)

ZAGRYATSKIY, I.V.; PROLOV, I.M.; KISELEV, S.M.

Drying enameled ware by combustion gases. Prom.energ. 11 no.8:  
19-21 Ag '56. (MLBA 9:11)  
(Enameled ware) (Drying apparatus)

KISELEV, S.M.

Using the products of natural gas combustion for heating air in  
air-heat screens and tributary systems. Gaz. prom. 7 no.8:27-29  
'62. (MIRA 17:10)

KISELEV, S.N.; IGRUNKOVSKIY, M.T.

Universal machine for drilling oil drains in piston grooves. Bul.  
tekh.-ekon.inform. no.5:16-18 '58. (MIRA 11:7)  
(Drilling and boring machinery)

KISELEV, S.N.

Lathe chucks with an electromechanical drive. Mashinostroitel'  
no. 1:45-47 Ja '66 (MIRA 19:1)

L 16571-66 EWT(d)/EWT(m) /EWP(w)/EWP(v)/T/EWP(t)/ETI/EWP(k) IJP(c) JD/HM/HW/EM  
ACC NR: AP6019432 SOURCE CODE: UR/0135/66/000/006/0034/0035 19  
AUTHOR: Kiselev, S. N. (Engineer); Pichugin, V. S. (Engineer) 18  
ORG: none 8  
TITLE: Down-hand and all-position inert gas-shielded arc welding of avial alloy pipe  
joints 16  
SOURCE: Svarochnoye proizvodstvo, no. 6, 1966, 34-35  
TOPIC TAGS: welding, shielded arc welding, aluminum alloy, magnesium containing alloy, silicon containing alloy, alloy pipe welding, weld property  
ABSTRACT: Annealed and aged pipes, 115x3.5 or 128x3.9 mm in diameter, from the avial-type aluminum alloy of the Al-Mg-Si system containing 0.90% Si and 0.65% Mn, were MIG welded with a consumable electrode either in argon atmosphere or in a mixture of argon with 60—70 He. X-ray inspection revealed that pipe joints welded in a fixed position had pores up to 0.6 mm in diameter and tungsten inclusions 0.2—0.3 mm in diameter, while down-hand welded joints, as a rule, had no defects. The welds made with the Ar-He mixture had a tensile strength of 19.7 kg/mm<sup>2</sup> and a bend angle of 86.3 deg; the corresponding figures for the welds made with argon were 17.3 kg/mm<sup>2</sup> and 59.5 deg. Welds made with the Ar-He mixture also had lower porosity. Generally, the strength of the gas-mixture welded joints was 70—80% of the strength of the base  
Card 1/2 UDC: 621.791.753.9:546.29.62.462:669.715

L 46571-66

ACC NR: AP6019432

metal, compared with 65—70% for argon welded joints. Also, softening of the heat-affected zone was much less pronounced in gas-mixture welds. The higher strength of the gas-mixture welded joints is explained by a lower specific heat input, and the greater bend angle, by the higher homogeneity of welded metal. Prolonged aging at 100C increased the weld tensile strength to 24.6—25 kg/mm<sup>2</sup>, but decreased the bend angle to 28—29.2 deg. In tests, all unaged joints failed in the heat-affected zone, but after aging at 100C for 1000 hr the joints failed predominantly along the weld metal. This is because the weld metal strength is lower than the strength of the aged base metal and also because of weld defects. Orig. art. has: 2 figures and 1 table. [ND]

SUB CODE: 13/ SUBM DATE: none/ ATD PRESS: 5028

Card 2/2

aum

L 27263-66 EWP(k)/EWT(d)/EWT(m)/EWP(h)/I/EWP(l)/EWP(v)/EWP(t) JD/HM

ACC NR: AP6009524

SOURCE CODE: UR/0413/66/000/005/0048/0048

AUTHORS: Kiselev, S. N.; Dedkov, L. K.; Schetchikov, B. A.; Pichugin, V. S.; Prosvirin, A. P.; Gamatudinov, B. I. 33  
B

ORG: none

TITLE: Automatic welder. Class 21, No. 179402 18

SOURCE: Izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, no. 5, 1966, 48

TOPIC TAGS: welder, butt welding, seam welding 14

ABSTRACT: This Author Certificate presents an automatic welder, using a nonmelting electrode in a protective atmosphere for ring and seam pipe welding. The welder includes an inlet port, ring-shaped rotator, welding head, system of roller supports, mechanisms for moving and correcting the welding head, electrode wire supplies, programmed current switching, and remote control equipment. To permit welding of variable diameter pipe and welding of flanges and rings, the rotator is equipped with a mechanism for displacement in the vertical plane, allowing a rotator body angle of 0--105° with respect to the horizontal. The centering mechanism consists of a fixture which is equipped with grips and shimming rings and a conical screw-driven compensator (see Fig. 1). A second feature has two perpendicular worms as

Card 1/2

UDC: 621.791.856.037

L 27263-66

ACC NR: AP6009524




Fig. 1. 1 - vertical displacement mechanism; 2 - centering mechanism; 3 - ring; 4 - conical compensators.

the rotator moving mechanism. Orig. art. has: 1 figure.

SUB CODE: 13/ SUBM DATE: 22Jun64/

Card 2/2 CC

ACC NR: AP7001836

(N)

SOURCE CODE: UR/0135/66/000/012/0006/0008

AUTHOR: Kiselev, S. N.; Khavanov, V. A. (Engineer); Skorniyakov, L. M. (Engineer); Grechishkin, V. I. (Engineer)

ORG: none

TITLE: Pattern of distribution of residual surface stresses in welded plates of avial alloy

SOURCE: Svarochnoye proizvodstvo, no. 12, 1966, 6-8

TOPIC TAGS: *welding equipment*  
metal stress, internal stress, weld evaluation, strain gage / Sv-AK-5 welding rod

ABSTRACT: The increasing use of avial-alloy-type structural elements and weldments of considerable thickness in which residual welding stresses combine with the scale factor as well as with the mechanical, chemical and structural heterogeneity of welded joints and the changes in plasticity of the material owing to aging processes, makes increasingly imperative an investigation of these stresses. Accordingly these stresses were measured in plates 30-, 40-, 70-, 90-, 140-, 220- and 300-mm thick of an avial type alloy containing 0.8-0.85% Si and 0.6-0.7% Mg in hardened and artificially aged state, with the aid of strain gages having a base of 5 mm and a resistance of the order of 50 ohm. The strain gauges were attached at intervals

Card 1/2

UDC: 621.791.011:669.715

ACC NR: AP7001836

of 100 mm each to the welded plates (which were 500 mm wide each half, and 500 and 1500 mm long) along the weld line in both directions from the center (in the direction of the principal axes of deformation). Findings: the pattern of distribution of residual surface stresses the welded joints of avial type plates differs from the pattern observed for low-carbon steels. Thus, in avial-type plates the residual welding stresses reach their maximum in the near-weld zone whereas in low-carbon steel plates these stresses reach their maximum at the weld center. This is attributable to the mechanical heterogeneity of the welded joints of avial-type alloys (the use of Sv-AK-5 welding rod, which contains 5% Si, and the softening of the base as well as to the features of formation of residual stresses, which are also determined by the thermophysical properties of the material: the high thermal conductivity of aluminum alloy leads to the elastic deformation of the metal in the near-weld zone. Orig. art. has: 5 figures.

SUB CODE: 13, 11, 20/ SUBM DATE: none/ ORIG REF: 003

Card 2/2

L 27380-66 EWT(m)/EWA(d)/EWP(v)/I/EWP(+)/ETI/EWP(k) IJP(c) YD/HM/JH  
 ACC NR: AP6015242 (A) SOURCE CODE: UR/0125/66/000/005/0016/0019  
 AUTHOR: Kiselev, S. N. (Moscow); Khovanov, V. A. (Moscow); Malyukov, V. A. (Moscow);  
 Skorniyakov, L. M. (Moscow); Matyunina, A. T. (Moscow)

ORG: none

TITLE: Mechanical properties of heavy welded avial-type alloy specimens

SOURCE: Avtomaticheskaya svarka, no. 5, 1966, 16-19

TOPIC TAGS: aluminum alloy, alloy weld, weld property, avial alloy

ABSTRACT: The effect of the size factor on the mechanical properties of heat-treatable avial-type aluminum-base alloy (0.74—0.90% Si, 0.59—0.70% Mg) welds and base metal has been studied. Specimens 10x10x100, 30x30x450, 40x40x500, 60x60x600, and 120x120x1000 mm (respective size factors 1, 3, 4, 6 and 12) were made from plates 40, 70, 90, 220 and 330 mm thick. Welding was done with a consumable SvAK-5 electrode in an argon-helium atmosphere. The base metal in the heat-treated condition (annealing and aging) had a tensile strength of 20—25 kg/mm<sup>2</sup>, a yield strength of 10—14 kg/mm<sup>2</sup>, and an elongation of 20—25%; corresponding figures for welded specimens were 16—19 kg/mm<sup>2</sup>, 8—10 kg/mm<sup>2</sup>, and 10—12%. Fracture in most cases was in the weld. Bend tests (on specimens with the Charpy-type notch) showed that with increasing size factor, the bend angle (measured at the appearance of the first crack)

Card 1/2

UDC: 621.791.053:620.172

L 27380-66

ACC NR: AP6015242




Fig. 1. Effect of the size factor on the bend angle of avial-type alloy base metal in the heat-treated condition (1) and aged at 130C for 200 hr (2); and in as-welded alloy welds (3) aged at 130C for 200 hr (4).

dropped. The values of bend angle in welded specimens were much lower than in base metal (see Fig. 1). Aging at 100 C for 1000 hr brings about a further drop in the ductility of welded specimens. After this treatment they failed in a brittle manner. Orig. art. has: 5 figures and 3 tables. [AZ]

SUB CODE: 11, 13/ SUBM DATE: 25May65/ ORIG REF: 005/ ATD PRESS: 4259

Card 2/2 20

25(2)

SOV/117-59-5-13/30

AUTHORS: Kiselev, S.N., Engineer, and Gribakin, M.N.

TITLE: A Hydraulic Device for the Scrap-Less Cold Rolling of Bar Stock for Making Keys

PERIODICAL: Mashinostroitel', 1959, Nr 5, pp 23-24 (USSR)

ABSTRACT: To cut down metal losses into chips, and the high machining costs in making keys, some plants are using cold rolling of bars in broaching machines. The rolled bars are then machined, so that there is still some machining and waste. The Vsesoyuznyy proyektno-tekhnologicheskii institut tyazhelogo mashinostroyeniya (All-Union Design and Technology Institute of Heavy Machine Building), jointly with the Mogilevskiy zavod pod'yemno-transportnogo oborudovaniya (Mogilev Materials Handling Machine Plant) have developed a special hydraulic device (Figure 2) for the "7510" broaching machine, with a 10-ton drawing effort. The exchangeable rollers of the device permit the obtaining of key bars of 8x10, 6x6 and 5x5 mm sections. The device is to be placed on the transition faceplate of the machine. The special chuck (Figure 3), fixed

Card 1/2

SOV/117-59-5-13/30

A Hydraulic Device for the Scrap-Less Cold Rolling of Bar Stock for Making Keys

in the machine slider, grips and pulls the rods in the rolling process. Design details are given. The application of the device at the Mogilev Plant has fully eliminated the forging and machining (planing and grinding) of keys, reduced by 25 to 30% the metal consumption, and improved the dimensions, accuracy and the mechanical properties of keys. There are 3 sets of diagrams.

Card 2/2

L 00996-66 EWT(d)/EPA(s)-2/EWT(m)/EWP(w)/EWA(d)/EWP(v)/T/EWP(t)/EWP(k)/EWP(z)/  
EWP(b)/EWA(c) JJP(c) EM/MJW/JD/HM  
ACCESSION NR: AP5018699 UR/0125/65/000/007/0044/0047  
621.791.856:669.715

AUTHOR: Kiselev, S. N. (Engineer) (Moscow); Khovanov, V. A. (Engineer) (Moscow);  
Skorniyakov, L. M. (Engineer) (Moscow); Malyukov, V. A. (Engineer) (Moscow)

TITLE: Welding thick plates of SAB-1 aluminum alloy

SOURCE: Avtomaticheskaya svarka, no. 7, 1965, 44-47

TOPIC TAGS: aluminum alloy, aluminum alloy thick plate, thick plate welding,  
edge groove geometry, welding electrode, weld metal property, heat treatment effect

ABSTRACT: Experiments have been made to develop an improved technique for welding thick plates of SAB-1 aluminum alloy, an age-hardenable alloy of the Al-Mg-Si system with Si:Mg > 1. Plates, 40, 80, and 140 mm thick, of SAB-1 alloy containing 0.81% Si and 0.48% Mg were inert-gas arc welded with a consumable electrode of the SvAK-5 type, 2, 4, or 5 mm in diameter, using a mixture of 30-40% Ar and 60-70% He for arc shielding. The use of helium made it possible to increase the temperature of the molten metal pool, to raise the voltage, and to ensure good weld formation. The best groove geometry was a double-V without root opening. In the experiments, the welding current was 450-520 amp, the arc voltage was 29-32 v, the Ar consumption

Card 1/2

L 00996-66

ACCESSION NR: AP5018699

was 30—35 l/min, and the He consumption was 50—60 l/min. The welding speed varied from 11.5 to 18.2 m/hr, and the number of passes was 6, 12—14, and 26—28 for plates 40, 80, and 140 mm, respectively. Welding with 4-mm electrode wire produced the least porous weld metal. Prior to heat treatment, the hardness of the heat-affected zone in 40-mm plates decreased by 15—18 HB compared with the parent metal, with the maximum decrease taking place at a distance of 12—15 mm from the fusion line. The corresponding figures for 80-mm plates were 10—12 HB and 8—10 mm, and for 140-mm plates, 5—8 HB and 5—6 mm. Subsequent heat treatment leveled to some extent the mechanical properties of the metal in the heat-affected zone, but did not improve them in the weld metal. Development of special electrode wire for welding SAB-1 type alloys is recommended to obtain welded joints which, after heat treatment, would have the strength of the parent metal. Orig. art. has: 5 figures and 3 tables.

[MS]

ASSOCIATION: none

SUBMITTED: 29Aug64

ENCL: 00

SUB CODE:MMIE

NO REF SOV: 004

OTHER: 001

ATD PRESS: 4068

Card 2/2

KISELEV, S.P.; KUDASOVA, G.F., kand. tekhn. nauk, red.; PANOV, A.A.,  
inzh., retsenzent; GLYASS, V.D., inzh., red.; LEYKINA, T.L.,  
red. izd-va; POL'SKAYA, R.G., tekhn. red.

[Metal polishing] Polirovanie metallov. Pod obskhei red. G.F.  
Kudasova. Moskva, Mashgiz, 1961. 67 p. (Bibliotekhka shlifov-  
shchika, no.10) (MIRA 14:12)

(Grinding and polishing)

YEVDOKIMOV, Boris Ivanovich; KISELEV, S.P., inzh.-podpolkovnik, red.;  
SOKOLOVA, G.F., tekhn.red.

[Rocket-powered antitank missiles; based on the foreign press]  
Protivotankovye upravlieniye reaktivnye snaryady; po materialam  
inostrannoi pechati. Moskva, Voen.izd-vo M-va obor.SSSR, 1959.  
73 p. (MIRA 13:3)

(Antitank guns)

(Rockets (Ordnance))

YEGOROV, Pavel Timofeyevich, kand.voyennykh nauk; KISELEV, S.P.,  
inzh.-podpolkovnik, red.; KONOVALOVA, Ye.K., tekhn.red.

[Rocket missiles] Reaktivnoe oruzhie. Moskva, Voen.isd-vo  
M-va obor.SSSR, 1960. 224 p. (MIRA 13:7)  
(Rockets (Ordnance))

VOLKOV, Yevgeniy Borisovich, dots., kand. tekhn. nauk, inzh.-polkovnik;  
KISELEV, S.P., red.; KATANUGIN, M.Ye., red.; KRASAVINA, A.M.,  
tekhn. red.

[Rocket engines] Raketnye dvigateli. Moskva, Voen. izd-vo M-va  
oborony SSSR, 1961. 58 p. (MIRA 14:12)  
(Rockets (Aeronautics))

VERMISHEV, Yuriy Khristoforovich, kand. tekhn. nauk, inzh.-polkovnik;  
PRIKHOD'KO, A.A., red.; KISELEV, S.P., red.; MEDNIKOVA, A.N.,  
tekhn. red.

[Rocket guidance] Upravlenie raketami. Moskva, Voen.izd-vo  
M-va obor.SSSR, 1961. 75 p. (MIRA 15:1)  
(Guided missiles) (Remote control)

MOROZOV, Pavel Vasil'yevich, inzh.-major; KATANUGIN, M.Ye., red.;  
KISELEV, S.P., red.; MEDNIKOVA, A.N., tekhn. red.

[Guided rocket weapons] Upravliaemoe raketnoe oruzhie. Moskva,  
Voen. izd-vo M-va oborony SSSR, 1961. 87 p. (MIRA 14:12)  
(Guided missiles)